

Astro 507  
Lecture 2  
Jan 24, 2014

Announcements:

- Preflight 1 due Fri. Jan 31, noon `www: assignment`

Note: answer in *two parts*

1. reading response: private, only I see
2. open-ended discussion question: public, everyone sees

Last time: Overview

at the **University** of Illinois we promise the whole universe

...it's right there in the name!

in this course: we deliver!

- └ Today's Agenda: The great work begins!
  - ★ cosmologist's observational toolbox
  - ★ zeroth-order structure of the Universe

## Program Notes: ASTR 507 Bugs/Features

- ▶ notes online—but come to class!  
if you print out, 4 pages/sheet works for me
- ▶ class  $\in$  diverse backgrounds: ask questions!  
also: occasionally need to be patient
- ▶ Socratic questions
- ▶ typos/sign errors  
Dirac story  
please report errors in lectures and problem sets;  
email notifications sent out

# Physical Cosmology

Modest goals:

scientific understanding of the

- origin
- evolution
- contents
- structure
- future

of the Universe

To be a science: must have empirical evidence

→ need observable data to reveal/test the above

ω *Q: What are cosmological observables?*

*hint: there are a wide variety*

# Cosmological Observables

“Raw” – hot off the instrument

## Local: Terrestrial/Solar System

- meteorites
- lunar samples
- solar wind

## Nonlocal: “Heavenly Messengers”

- photon signals: individual objects
  - local and Galactic: Sun, stars, gas
  - extragalactic: galaxies, QSO, etc
- diffuse photon backgrounds (all  $\lambda$ : radio–gamma ray)
- cosmic rays
- neutrinos
- gravity waves (!)
- dark matter particles (?)

## “Cooked” – After Analysis

- *composition* of meteorites, solar wind, moonrocks  
elements, isotopes
- photon spectra of stars, galaxies, interstellar/intergalactic gas  
→ element composition, red/blueshifts, temperature
- galaxy distribution
- galaxy distortions (lensing)
- masses, velocities, radii, ...

Armed with these, we proceed...

# Bizarre Astronomical Units I: Distances

Charity begins at home: *Astronomical Unit (AU)*

- average Earth-Sun distance (really: semi-major axis); known very precisely
- $a(\text{Earth} - \odot) \equiv 1 \text{ AU} = 1.49597870660 \times 10^{13} \text{ cm}$


## parsec

- star with parallax angle  $p$  lies at distance

$$r(p) = \frac{1 \text{ AU}}{\tan p} \approx \frac{1 \text{ AU}}{p}$$

- for  $p = 1 \text{ arcsec} = 4.8 \times 10^{-6} \text{ rad}$ , distance is

○  $r(1 \text{ arcsec}) \equiv 1 \text{ parsec} \equiv 1 \text{ pc} = 3.0857 \times 10^{18} \text{ cm} \approx 3 \text{ yr} \quad (1)$




## Cosmologist's Toolbox: Energy Flow

idealized detector of area  $dA$

receives all incident radiation (all directions, all  $\nu$ )

over exposure time  $dt$



energy received in exposure  $d\mathcal{E}$  depends on detector

because  $d\mathcal{E} \propto dA dt$  Q: why?

thus energy received is detector-dependent via  $dA$

Q: how to remove detector dependence?

## Energy Flux and Inverse Square Law

independent of detector, and  
intrinsic to source and distance: **energy flux** (or just “flux”)

$$F = \frac{dE}{dA dt} = \frac{d\text{Power}}{d\text{Area}} \quad (2)$$

cgs units:  $[F] = [\text{erg cm}^{-2} \text{s}^{-1}]$


consider stationary spherical source of size  $R$   
in a non-expanding, Euclidean space  
emitting isotropically = uniformly in all directions  
with constant power  $L$  (“luminosity”)

at radius  $r > R$  (outside of source)  
area  $A = 4\pi r^2$ , and flux is

$\infty$

$$F = \frac{L}{4\pi r^2}$$

*inverse square law*





# The Shape and Scale of the Universe

basic & ancient questions:

- how old is the universe?
- how big is the universe?
- what is the universe made of?

note: one's idea of "universe" implicitly presupposes aspects of answers to these questions

historically:

- dramatic upward revisions in scale of U
- drastic broadening in known cosmic composition

from Newton to early 20th Century:

- $\Rightarrow$  *Universe*  $\equiv$  *Milky Way*: a (finite) collection of *stars*  
size  $\sim$  *few* kpc = *few*  $\times 10^3$  pc

# The Realm of the Nebulae

*one century ago...*

hottest question in 1920's astronomy:

what are spiral "nebulae"?

www: Milky Way and Galactic coordinates

www: NGC sky and Zone of Avoidance

tool: Cepheids—variable stars

www: Cepheids then and now

⇒ periodically changing **luminosity**

*Q: how to measure  $P$ ?  $L$ ?*

Henrietta Leavitt & Charles Pickering (1912):  $P - L$  correlation  
in LMC stars

⊕ calibrate  $L$  vs  $P$  relation locally, then apply to nebulae:

from period  $P$  ⇒ infer  $L$

If lucky or clever:  $L$  known ( “**standard candle**” )  
solve for “**luminosity distance**”

$$d_L = \sqrt{\frac{L}{4\pi F}} \quad (3)$$

Edwin Hubble: Cepheids in “Andromeda Nebula” M31

$$d_L \sim 10^3 \text{ kpc} \gg R_{\text{MilkyWay}}$$

⇒ M31 is “island universe” = galaxy

cosmic distance scale grew by factor  $\sim 1000$ : kpc → Mpc

So to summarize:

Q: pc, kpc, Mpc, Gpc *characteristic scales for what?*

## Typical Lengthscales: Cosmic Hierarchy

- ★ typical **star-star separation** in galaxies  $\sim 1$  pc
- ★ typical (visible) **galaxy size**  $\sim 1\text{kpc} = 10^3$  pc
- ★ (present-day) typical **galaxy-galaxy separation**  
 $\sim 1$  Mpc  $= 10^6$  pc
- ★ (present-day) **observable universe**  $\sim 1$  Gpc  $= 10^9$  pc

Q: *Why is this a "hierarchy"?*

# Observational Cosmology: Zeroth-Order Picture

## Cosmic Matter Distribution

*Q: how quantify distribution?*

*Q: how characterize smoothness/lumpiness?*

*Q: how determine observationally?*

# Galaxy Maps and Cosmic Structure

observable cosmo “building blocks” – galaxies  
 $\approx$  all stars in galaxies

www: Galaxy Survey: 2dFGRS

map galaxies in “slices” of sky  $2^\circ$  thick

*Q: qualitative trends—small scales? large scales?*

*Q: how could we make this more quantitative?*

*Q: how to test these conclusions?*